

Journal Pre-proofs

The number of leisure-time activities and risk of functional disability among Japanese older population: the JAGES cohort

Yoshihiro Fuji, Ryoto Sakaniwa, Kokoro Shirai, Tami Saito, Shigekazu Ukawa, Hiroyasu Iso, Katsunori Kondo

PII: S2211-3355(22)00048-1
DOI: <https://doi.org/10.1016/j.pmedr.2022.101741>
Reference: PMEDR 101741

To appear in: *Preventive Medicine Reports*

Received Date: 8 September 2021
Revised Date: 14 February 2022
Accepted Date: 16 February 2022



Please cite this article as: Y. Fuji, R. Sakaniwa, K. Shirai, T. Saito, S. Ukawa, H. Iso, K. Kondo, The number of leisure-time activities and risk of functional disability among Japanese older population: the JAGES cohort, *Preventive Medicine Reports* (2022), doi: <https://doi.org/10.1016/j.pmedr.2022.101741>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The number of leisure-time activities and risk of functional disability among Japanese older population: the JAGES cohort

Abstract

Evidence-based prevention of functional disability is a pressing issue for the health among the older population, due to the rapidly global aging. This study aimed to examine the association between leisure-time activities and the risk of functional disability. In a longitudinal prospective cohort study, we recruited 50,286 Japanese men and women aged ≥ 65 years who did not have functional disability at the baseline in 2010–2011, with a median follow-up of 5.8 years. We examined the association between 24 leisure-time activities and the risk of developing functional disability. Cox proportional hazards regressions were used to examine the hazard ratios (HRs) and 95% confidence intervals (95% CIs) for functional disability. A total of 10,631 persons (4,497 men and 6,134 women) newly developed functional disability. The number of leisure-time activities was inversely associated with the risk of functional disability. With reference to no activity, the multivariable HRs (95% CIs) were 0.89 (0.82–0.97) for one activity, 0.72 (0.67–0.78) for two to four activities, and 0.66 (0.58–0.74) for five or more activities (P for trend, <0.001) for men, and for women, the corresponding HRs were 0.84 (0.78–0.90), 0.77 (0.72–0.82), and 0.70 (0.62–0.79), (P for trend, <0.001). Further, even lower-loading leisure-time activities such as computer for men and handicrafts for women, were also associated with a reduced risk of functional disability. Our study suggests the importance of engaging in various leisure-time activities among the older population.

Keywords

leisure-time activities, functional disability, older population, lower-loading, prospective study

Yoshihiro Fuji, MPH, jzx90.mark23669@gmail.com^{a,b}

Ryoto Sakaniwa, MPH, PhD, r.sakaniwa@pbhel.med.osaka-u.ac.jp^a

Kokoro Shirai, MPH, PhD, kshirai@pbhel.med.osaka-u.ac.jp^a

Tami Saito, PhD, t-saito@ncgg.go.jp^c

Shigekazu Ukawa, PhD, ukawa@life.osaka-cu.ac.jp^d

Hiroyasu Iso, MD, MPH, PhD, iso@pbhel.med.osaka-u.ac.jp^a

Katsunori Kondo, MD, PhD, kkondo@kkondo.net^{e,f}

^a Public Health, Department of Social Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan, 2-2 Yamadaoka, Suita-shi, Osaka, 565-0871 Japan

^b Department of Rehabilitation, Osaka General Medical Center, Osaka, Japan, 3-1-56, Sumiyoshi-ku, Osaka-shi, Osaka, 558-8558, Japan

^c Department of Social Science, Center for Gerontology and Social Science, National Center for Geriatrics and Gerontology, Aichi, Japan, 7-430, Morioka-cho, obu-shi, Aichi, 474-8511, Japan

^d Osaka City University Graduate School of Human Life Science, Osaka, Japan, 3-3-138, Sugimoto, Sumiyoshi-ku, Osaka-shi, Osaka, 558-8585, Japan

^e Department of Social Preventive Medical Sciences, Center for Preventive Medical Sciences, Chiba University, Chiba, Japan, 1-8-1, Inohana, Chuo-ku, Chiba-shi, Chiba, 260-0856, Japan

^f Department of Gerontological Evaluation, Center for Gerontology and Social Science, National Center for Geriatrics and Gerontology, Aichi, Japan, 7-430, Morioka-cho, obu-shi, Aichi, 474-8511, Japan

Functional disability are not a natural consequence of aging and could be prevented in primary healthcare settings (Yates and Muchisky, 1997; WHO, 2008). Evidence-based prevention of functional disability is a pressing issue for the health among the older population, particularly due to the rapidly aging population in several Asian countries (WHO, 2017; eurostat Statistics Explained). In Japan, the annual medical costs for managing patients with functional disability are expected to increase from US \$ 100 billion in 2018 to US \$ 150 billion by 2025 (Labour and Welfare. Ministry of Health; Labour and Welfare. Ministry of Health). Worldwide, the number of years lived with disability doubled from 17.7 million in 1990 to 34.4 million in 2019 (Vos et al., 2020).

The association of leisure-time activities and health benefits among the older population has been described based on the levels of physical activity (Corbett et al., 2018; Chen et al., 2016). For instance, within a median follow-up of 11.6 years, the New Integrated Suburban Seniority Investigation (NISSIN) of 2,888 Japanese men aged ≥ 65 years showed that leisure-time activity with ≥ 18 metabolic equivalent (MET) hours/week of leisure-time activities was associated with a reduced risk of functional disability compared with no leisure-time activities (multivariable hazard ratio [HR]: 0.48; 95% confidence interval [CI]: 0.25–0.94) (Matsunaga et al., 2017). A 10-year follow-up prospective study of 2,456 Finnish men and women aged ≥ 65 years showed that participation in intensive sports and training for ≥ 3 hours/week were associated with a reduced risk of incident cardiovascular disease (coronary heart disease and stroke) compared with non-intensive leisure-time activities (HR: 0.55, 95% CI: 0.38–0.79) (Barengo et al., 2017).

The Bronx aging study of 469 English men and women aged ≥ 75 years reported that many of the older adults participated two or more types of leisure-time activities such as playing board games (22.6%), reading books (81.4%), writing (18.6%), performing housework (77.4%), walking (86.1%), climbing the stairs (67.4%), and participating in group exercise (30.0%) (Verghese et al., 2003). Therefore, the impact of the number of various leisure-time activities on the risk of functional disability should be taken into account.

Previous studies reported the benefit of the number of leisure-time activities on the risk of all-cause mortality (Kobayashi et al., 2021), dementia (Xu et al., 2017; Ling et al., 2020), and cognitive impairments

number and frequency of social participations in leisure-time activities were associated with a lower risk of functional disability (Kanamori et al., 2014; Komatsu et al., 2019). However, the association between the number of different types of leisure-time activities such as walking/running, and gardening, traveling and the risk of functional disability has not been investigated.

This prospective cohort study aimed to examine the association between the number of various leisure-time activities and the risk of functional disability in 50,286 Japanese individuals aged ≥ 65 years.

Methods

Study population

We used the data from the JAGES (Japan Gerontological Evaluation Study) cohort. The study profile has been described elsewhere (Kondo, 2016; Kondo K et al., 2018). In brief, the baseline mail survey was conducted in 2010–2011 using a self-administered questionnaire. A total of 54,539 people (25,146 men and 29,393 women) aged ≥ 65 years without functional disability from 31 municipalities in Japan were enrolled in our study. We excluded 48 individuals who did not respond to our survey and 4,204 individuals with missing data on leisure-time activities. Finally, 50,286 (23,103 men and 27,183 women) individuals were included in the current study. Data on the incidence of functional disability from 2010 to 2016 were obtained from municipal public long-term care insurance (LTCI) system databases, and we identified people who were newly eligible for the LTCI benefit considered as those with functional disability.

This study was ethically approved by the institutional review boards (no. 10-05, no. 1777). All respondents were informed that participation in this study was absolutely voluntary and that completing and returning the questionnaires via mail indicated their consent to participate.

Definition for functional disability

The follow-up period started from August 2010 until April 2016, with the longest being 2,070 days. Older adults in Japan with functional disability were certified for LTCI database (Labor and Welfare Ministry of

functional disability (Tsutsui and Muramatsu, 2005; Tsutsui and Muramatsu, 2007). This system was applied as public and social welfare services throughout Japan and covers aged ≥ 65 years with limitations in performing ADL and/or instrumental ADL (IADL) regardless of their income status. The levels of functional disability were assessed by certified interviewers through home-visits and based on written opinions from a primary physician, including a physician's examination and evaluation of physical and cognitive functions (Tsutsui and Muramatsu, 2005). Functional disabilities were assessed with seven levels: support needed (levels 1 to 2), care needed (levels 1 to 5), or no functional disability (independent). All registered individuals who were identified with functional disability received public and social welfare services for adult day care, home-visit care, day-visit services, short-stay services, residential services, and/or in-facility services, depending on the degree of functional disabilities from local government (Labour and Welfare Ministry of Health). These LTCI's criteria are widely adopted in previous epidemiological studies (Kanamori et al., 2014; Ukawa et al., 2020; Hikichi et al., 2015; Ashida et al., 2016; Aida et al., 2013). In each, local Japanese government, the applicants for the LTCI are asked to fill out a basic application form (Ministry of Health, 2016).

Information on all-cause mortality was obtained from the LTCI database system of the participants' municipality office. Those who died from any cause, moved out from their original community, or had no data on the last follow-up, whichever came first, were censored.

Definition of the number of leisure-time activities

At baseline, all participants were asked the following question: "Do you currently have any leisure-time activities?" When the participants responded "yes," they were asked to provide the different types of leisure-time activities that they engaged in.

According to physical activities tracking guide, the amount of time spent (in METs) in performing leisure-time activities were as follows: for example walking/running: 2.3/6.0, gardening: 3.3, traveling: 2.5, reading: 1.3, computer: 1.5, and handicrafts: 1.3 (Ainsworth BE et al.,). The types of leisure-time activities were classified as: higher-loading leisure-time activities (METs \geq approximately 2.5: walking/running,

exercise/tai chi, mountain climbing, instrument performance, gate ball, dance, and chorus/folk song) and lower-loading activities (METs <2.5: reading, computer, igo/shogi/mahjong, pachinko, painting/picture letter, calligraphy, haiku/tanka/senryu, handicrafts, and tea ceremony/flower arrangement), and other non-specified activities.

We counted the total number of leisure-time activities performed by each individual (from 0 to 24 points). All participants were divided into four groups according to the number of leisure-time activities (zero, one, two to four, and \geq five or more types).

Covariates

The sociodemographic variables (sex, age, and marital status), socioeconomic status (educational level, occupational status, and equivalized income), health-related behaviors (smoking status, IADL status, and walking hours), and histories of comorbidities (diabetes mellitus, hypertension, stroke, heart disease, mental disorder, and hearing loss), engagement in social activities (political groups/organizations, volunteer groups, sports groups/clubs, neighborhood associations, and senior clubs), and frequency of meeting friends were included as covariates for multivariate-adjusted examinations. These variables were divided into the following categories: sex (men or women), age (65–69, 70–74, 75–79, 80–84, or \geq 85 years), educational level (<10 or \geq 10 years), occupational status (employed or not employed), equivalized income (<200, 200–399, or \geq 400 \times 10,000 JPY; 100 JPY \approx 1 USD), marital status (married, widowed, divorced, or never married), smoking status (yes or no), IADL status (independent; 5 points out of 5 or not-independent; less than 5 points) (Koyano et al., 1991), walking hours (<30, 30–59, 60–89, \geq 90 min/day), and histories of comorbidities (diabetes mellitus, hypertension, stroke, heart diseases, mental disorder, and/or hearing loss), social activities (political groups/organizations, volunteer groups, sports groups/clubs, neighborhood associations, and/or senior clubs), and frequency of meeting friends (\geq four or more times/week, two to three times/week, one time/week, one to two times/month, less than one times/year).

Statistical Analyses

examined using the chi-square test for categorical variables, one-way analysis of variance (ANOVA) for continuous variables, and Cochran-Armitage test for variables expressed as percentage. Cox proportional hazards regression was used to calculate the HR and 95% CI of functional disability after controlling for confounding variables. We also calculated the HR of functional disability associated with higher- and lower-loading leisure-time activities, separately. The confounding variables for the multivariable adjustment in model 1 were age, educational level, occupational status, equivalized income, marital status, smoking status, IADL status, walking hours, histories of diabetes mellitus, hypertension, stroke, heart disease, mental disorder, and hearing loss. Model 2 was adjusted further for engagement in social activities of political groups/organizations, volunteer groups, sports groups/clubs, neighborhood associations, senior clubs, and frequency of meeting friends. For participants with missing data, we imputed “missing-variable” as covariate. All missing data were considered as missing completely at random, and the number of participants with missing data were generally small. Meanwhile, functional disability that occurred from year 1 to year 3 were included in the sensitive analysis to reduce the possibility of reverse causation. *P*-values ≤ 0.05 (two-sided tails) were considered as significant. All statistical analyses were performed using SAS version 9.4 (SAS Institute Inc., Cary, NC, USA).

Results

The participant’s mean ages at baseline were 73.9 years for men and 74.2 years for women. The number of newly certified LTCI according to the number of leisure-time activities and sex were as follows: 1,643 for no activity, 933 for one activity, 1,522 for two to four activities, and 399 for five or more activities in men, and 2,712, 1,189, 1,865, 368, respectively in women.

Table 1 shows the sex-specific baseline characteristics according to the number of leisure-time activities. In brief, younger individuals, higher educational level, unemployed status, high IADL status, higher current equivalized income, ≥ 30 walking hours, the participation in political groups/organizations, volunteer groups, sports groups/clubs, neighborhood associations, senior clubs, and frequency of meeting friends were more prevalent in individuals with higher number of leisure-time activities for both men and

mental disorder, and hearing loss were inversely associated with the number of leisure-time activities for both men and women.

The proportion of each leisure-time activity according to the number of leisure-time activities is summarized in Table 2. Many of the participants performed walking/running (6,119 men, 4,936 women), gardening (5,675 men, 7,653 women), traveling (5,354 men, 5,742 women), reading (3,865 men, 3,805 women), computer (3,148 men, 1,002 women), and handicrafts (306 men, 3,782 women) as leisure-time activities.

During a median follow-up of 5.8 years, 10,631 participants (4,497 men, 6,134 women) developed functional disability. The sex-specific HRs and 95% CIs for functional disability, according to the number of leisure-time activities, are summarized in Table 3. The number of leisure-time activities was inversely associated with the risk of functional disability for both men and women even after adjustment for potential confounding variables. In the final model (model 2), the multivariable HRs (95% CIs) for functional disability were as follows: 0.89 (0.82–0.97) for one activity, 0.72 (0.67–0.78) for two to four activities, and 0.66 (0.58–0.74) for five or more activities (P for trend, <0.001) in men, 0.84 (0.78–0.90), 0.77 (0.72–0.82), 0.70 (0.62–0.79) (P for trend, <0.001), respectively in women. The dose-response associations were not attenuated, after exclusion of early onsets of functional disability from 1 to 3 years for both men and women (P for trend, <0.001).

The HRs and 95% CIs for functional disability, according to higher- and lower- loading leisure-time activities are summarized in Table 4. The higher loading leisure-time activities were inversely associated with the risk of functional disability even after adjustment for potential confounding variables. The lower loading leisure-time activities also tended to be inversely associated with the risk of functional disability.

The sex-specific associations between each leisure-time activities and the risk of functional disability are shown in Supplementary Table 1. For both men and women, traveling, cultivation of agricultural crops were inversely associated with the risk of functional disability. An inverse association was also observed between walking/running, computer, fishing, golf, mountain climbing and the risk of

risk of functional disability for women, but no significant association was observed between other leisure-time activities and the risk for both men and women.

Discussion

In this large prospective cohort study of older men and women, we found inverse associations between the number of leisure-time activities and the risk of functional disability for both men and women. Compared with persons without leisure-time activities, those with one or more leisure-time activities had reduced risk of functional disability in a dose-response manner. These associations did not substantially change after exclusion of early onsets of functional disability.

In a previous JAGES study conducted in 13,310 Japanese aged ≥ 65 years, social participation in three or more in hobby clubs, sports groups, and local community activities were associated with a 43% lower risk of functional disability compared with no social participation in the 4-year follow-up (Kanamori et al., 2014). In another JAGES study conducted in 44,978 Japanese aged ≥ 65 years, the high frequency of social participation in both sports and the hobby and high frequency of social participation in both political and industry/trade activities were associated with reduced risks of functional disability by 34% and 29%, respectively compared with the low frequency of social participation in the 3-year follow-up (Ukawa et al., 2020). In a previous 3-year follow-up study of 6,360 Japanese older adults, the participation in hobby clubs was associated with a 32% reduced risk of functional disability for men and a 47% reduced risk of functional disability for women compared with no social participation (Tomioka et al., 2017). These previous studies, however, did not examine the association between the number of leisure-time activities and risk of functional disability.

We found an association between functional disability and various leisure-time activities such as traveling, cultivation of agricultural crops, walking/running (men only), computer (men only), fishing (men only), golf (men only), mountain climbing (men only), and gardening (women only), instrument performance (women only), dance (women only), handicrafts (women only). The guidelines by the Ministry of Health and Labour and Welfare, Japan recommend that older adults (aged ≥ 65 years) should perform at

years showed that persons who performed ≥ 18 MET-hours/weeks of leisure-time activities had a 52% lower risk of developing functional disability, compared with those who did not engage in leisure-time activities during 11.6 years median follow-up (Matsunaga et al., 2017). By contrast, our study showed that even lower-loading leisure-time activities tended to be associated with a reduced the risk of functional disability. We assume that lower-loading activities likely make older people to be less physically fatigued, and to be continued as habits. Most of lower-loading activities needs manual dexterity which was associated with a lower risk of mild cognitive impairment in 7-year follow-up of 1,160 older Americans (Beeri M.S., 2021).

Our study has two strengths. First, we followed up over 50,000 older adults for a median of 5.8 years. Second, the 24 types of leisure-time activities were tested to examine the impact of the number of leisure-time activities on functional disability.

This study has several limitations. First, objective assessment for clinical diagnosis of functional disability was not conducted systematically at enrollment so that in some participants functional disability might have existed in our study. Second, the reverse causation for the association between leisure-time activities and the risk of functional disability remained. However, the absence of substantial changes in the association after the exclusion of early onsets of functional disability from the baseline up to 3 years suggested that the reverse causation may be unlikely.

Conclusion

We found inverse dose-response associations between the number of leisure-time activities including lower-loading activities and the risk of functional disability among Japanese older men and women. Our finding suggests that engaging in various types of leisure-time activities may contribute to the prevention of functional disability.

Declaration of competing interest

None.

We do not have the contribution of others who merit authorship and sponsor.

Funding

This study used data from JAGES (The Japan Gerontological Evaluation Study). This study was supported by Grants-in-Aid for Scientific Research (15H01972, 15H04781, 15H05059, 15K03417, 15K03982, 15K16181, 15K17232, 15K18174, 15K19241, 15K21266, 15KT0007, 15KT0097, 16H05556, 16K09122, 16K00913, 16K02025, 16K12964, 16K13443, 16K16295, 16K16595, 16K16633, 16K17256, 16K17281, 16K19247, 16K19267, 16K21461, 16K21465, 16KT0014, 16KK0059, 17K04305, 17K34567, 17K04306, 25253052, 25713027, 26285138, 26460828, 26780328, 18H03018, 18H04071, 18H03047, 18H00953, 18H00955, 18KK0057, 19H03901, 19H03915, 19H03860, 19K04785, 19K10641, 19K11657, 19K19818, 19K19455, 19K24060, 19K20909, 20H00557) from JSPS (Japan Society for the Promotion of Science); Health Labour Sciences Research Grants (h26-Choju-Ippan-006, h27-Ninchisyou-Ippan-001, h28-Choju-Ippan-002, h28-Ninchisyou-Ippan-002, h29-Chikyukibo-Ippan-001, h30-Jyunkankinado-Ippan-004, 18H04071, 19FA1012, 19FA2001) from the Ministry of Health, Labour and Welfare, Japan; the Research and Development Grants for Longevity Science from Japan Agency for Medical Research and development (AMED) (JP17dk0110027, JP18dk0110027, JP18ls0110002, JP18le0110009, JP20dk0110034, JP20dk0110037, 21lk0310073h0002), the Research Funding for Longevity Sciences from National Center for Geriatrics and Gerontology (20-19, 21-20, 24-17, 24-23, 29-42, 30-30,30-22); Open Innovation Platform with Enterprises, Research Institute and Academia (OPERA, JPMJOP1831) from the Japan Science and Technology (JST); a grant from the Japan Foundation For Aging And Health (J09KF00804), a grant from Innovative Research Program on Suicide Countermeasures (1-4), a grant from Sasakawa Sports Foundation, a grant from Japan Health Promotion & Fitness Foundation, a grant from Chiba Foundation for Health Promotion & Disease Prevention, the 8020 Research Grant for fiscal 2019 from the 8020 Promotion Foundation (adopted number: 19-2-06), a grant form Niimi University (1915010), grants from Meiji Yasuda Life Foundation of Health and Welfare, Osaka University Center of Medical Data Science Advanced Clinical Epidemiology Investigator's Research, and the OU-UCL Joint-Lab Research Center in Osaka

reflect the official policy or position of the respective funding organizations.

Journal Pre-proofs

- Aida, J., Kondo, K., Kawachi, I., et al., 2013. Does social capital affect the incidence of functional disability in older Japanese? A prospective population-based cohort study. *J. Epidemiol. Community Health* 67, 42–47. <https://doi.org/10.1136/jech-2011-200307>
- Ainsworth BE, Haskell WL, Herrmann SD, et al, Healthy Lifestyles Research Center, College of Nursing & Health Innovation, Arizona State University: The Compendium of Physical Activities Tracking Guide. <https://sites.google.com/site/compendiumofphysicalactivities/home> (accessed 9.1.2021).
- Ashida, T., Kondo, N., Kondo, K., 2016. Social participation and the onset of functional disability by socioeconomic status and activity type: The JAGES cohort study. *Prev. Med. (Baltim)*. 89, 121–128. <https://doi.org/10.1016/j.ypmed.2016.05.006>
- Beeri, M.S., Leurgans, S.E., Bennett, D.A., et al., 2021. Diverse Motor Performances Are Related to Incident Cognitive Impairment in Community-Dwelling Older Adults. *Front. Aging Neurosci.* 13. <https://doi.org/10.3389/fnagi.2021.717139>
- Barengo, N.C., Antikainen, R., Borodulin, K., et al., 2017. Leisure-Time Physical Activity Reduces Total and Cardiovascular Mortality and Cardiovascular Disease Incidence in Older Adults. *J. Am. Geriatr. Soc.* 65, 504–510. <https://doi.org/10.1111/jgs.14694>
- Chen, Y.M., Chiang, T.L., Chen, D.R., et al., 2016. Trajectories of Older Adults' Leisure Time Activity and Functional Disability: a 12-Year Follow-Up. *Int. J. Behav. Med.* 23, 697–706. <https://doi.org/10.1007/s12529-016-9554-y>
- Corbett, D.B., Rejeski, W.J., Tudor-Locke, C., et al., 2018. Social Participation Modifies the Effect of a Structured Physical Activity Program on Major Mobility Disability among Older Adults: Results from the LIFE Study. *Journals Gerontol. - Ser. B Psychol. Sci. Soc. Sci.* 73, 1501–1513. <https://doi.org/10.1093/geronb/gbx051>
- Doi, T., Verghese, J., Makizako, H., et al., 2017. Effects of Cognitive Leisure Activity on Cognition in Mild Cognitive Impairment: Results of a Randomized Controlled Trial. *J. Am. Med. Dir. Assoc.* 18, 686–691. <https://doi.org/10.1016/j.jamda.2017.02.013>

- https://ec.europa.eu/eurostat/statistics-explained/index.php/Functional_and_activity_limitations_statistics#Self-reported_long-standing_limitations_due_to_health_problems. (accessed 9.1.2021).
- Hikichi, H., Kondo, N., Kondo, K., et al., 2015. Effect of a community intervention programme promoting social interactions on functional disability prevention for older adults: Propensity score matching and instrumental variable analyses, JAGES Taketoyo study. *J. Epidemiol. Community Health* 69, 905–910. <https://doi.org/10.1136/jech-2014-205345>
- Kanamori, S., Kai, Y., Aida, J., et al., 2014. Social participation and the prevention of functional disability in older Japanese: The JAGES cohort study. *PLoS One* 9. <https://doi.org/10.1371/journal.pone.0099638>
- Kobayashi, T., Tani, Y., Kino, S., et al., 2021. Prospective Study of Engagement in Leisure Activities and All-Cause Mortality Among Older Japanese Adults. *J. Epidemiol.* <https://doi.org/10.2188/jea.je20200427>
- Komatsu, M., Obayashi, K., Tomioka, K., et al., 2019. The interaction effect between physical and cultural leisure activities on the subsequent decline of instrumental ADL: The Fujiwara-kyo study. *Environ. Health Prev. Med.* 24. <https://doi.org/10.1186/s12199-019-0826-4>
- Kondo, K., 2016. Progress in aging epidemiology in Japan: The JAGES project. *J. Epidemiol.* 26, 331–336. <https://doi.org/10.2188/jea.JE20160093>
- Kondo K, Rosenberg M., editors, 2018. Advancing universal health coverage through knowledge translation for healthy ageing. <https://apps.who.int/iris/handle/10665/279010> (accessed 9.1.2021).
- Ling, L., Tsuji, T., Nagamine, Y., et al., 2020. Types and number of hobbies and incidence of dementia among older adults: A six-year longitudinal study from the Japan Gerontological Evaluation Study (JAGES). *Nihon. Kosshu Eisei Zasshi.* 67, 800–810. https://doi.org/10.11236/jph.67.11_800, (in Japanese).
- Matsunaga, T., Naito, M., Wakai, K., et al., 2017. Leisure-time physical activity and risk of disability incidence: A 12-year prospective cohort study among young elderly of the same age at baseline. *J. Epidemiol.* 27, 538–545.

Ministry of Health, Labour and Welfare. Long-Term Care Insurance System of Japan.

https://www.mhlw.go.jp/english/policy/care-welfare/care-welfare-elderly/dl/ltcisj_e.pdf.

http://www.mhlw.go.jp/english/policy/care-welfare/care-welfare-elderly/dl/ltcisj_e.pdf (accessed 9.1.2021).

Ministry of Health, Labour and Welfare. Long term care insurance business status report (annual report)

Ministry of Health, Labour and Welfare.

<https://www.mhlw.go.jp/topics/kaigo/osirase/jigyo/18/index.html> (accessed 9.1.2021).

Ministry of Health, Labour and Welfare. The future outlook for social security with an eye on 2040 Ministry

of Health, Labour and Welfare. <https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000207382.html> (accessed 9.1.2021).

Ministry of Health, Labor and Welfare. Promotion of exercise measures.

https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/kenkou_iryoku/kenkou/undou/index.html (accessed 9.1.2021).

Ministry of Health, Labor and Welfare. Physical activity criteria to promote health 2013 [PDF on the Internet].

<http://www.mhlw.go.jp/stf/houdou/2r9852000002xple-att/2r9852000002xpqt.pdf> (accessed 9.1.2021).

Sugita, A., Ling, L., Tsuji, T., et al., 2020. Cultural engagement and incidence of cognitive impairment: A six-year longitudinal follow-up of the Japan Gerontological Evaluation Study (JAGES). *J. Epidemiol.* <https://doi.org/10.2188/jea.je20190337>

Tomioka, K., Kurumatani, N., Hosoi, H., 2017. Association Between Social Participation and 3-Year Change in Instrumental Activities of Daily Living in Community-Dwelling Elderly Adults. *J. Am. Geriatr. Soc.* 65, 107–113. <https://doi.org/10.1111/jgs.14447>

Tsutsui, T., Muramatsu, N., 2007. Japan's universal long-term care system reform of 2005: Containing costs and realizing a vision. *J. Am. Geriatr. Soc.* 55, 1458–1463. <https://doi.org/10.1111/j.1532-5415.2007.01281.x>

- J. Am. Geriatr. Soc. <https://doi.org/10.1111/j.1532-5415.2005.53175.x>
- Ukawa, S., Tamakoshi, A., Okada, Y., et al., 2020. Social participation patterns and the incidence of functional disability: The Japan Gerontological Evaluation Study. *Geriatr. Gerontol. Int.* 20, 765–772. <https://doi.org/10.1111/ggi.13966>
- Verghese, J., Lipton, R.B., Katz, M.J., et al., 2003. Leisure Activities and the Risk of Dementia in the Elderly. *N. Engl. J. Med.* 348, 2508–2516. <https://doi.org/10.1056/NEJMoa022252>
- Vos, T., Lim, S.S., Abbafati, C., et al., 2020. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 396, 1204–1222. [https://doi.org/10.1016/S0140-6736\(20\)30925-9](https://doi.org/10.1016/S0140-6736(20)30925-9)
- WHO, 2008. Age-Friendly Primary Health Care Centres Toolkit. http://apps.who.int/iris/bitstream/handle/10665/43860/9789241596480_eng.pdf;jsessionid=F18E6E06F7E0556DD7757F99DCF14F16?sequence=1 (accessed 9.1.2021).
- WHO, 2017. Global action plan on the public health response to dementia 2017 - 2025. https://www.who.int/mental_health/neurology/dementia/action_plan_2017_2025/en/ (accessed 9.1.2021).
- Xu, W., Wang, H.F., Wan, Y., et al., 2017. Leisure time physical activity and dementia risk: A dose-response meta-analysis of prospective studies. *BMJ Open* 7, 1–10. <https://doi.org/10.1136/bmjopen-2016-014706>
- Yates, R., Muchisky, D., 1997. Successful aging. *Gerontologist* 37, 433–440. <https://doi.org/10.5054/tq.2010.215250>

Table 1. Sex-specific baseline characteristics according to the number of leisure-time activities

		Men				Women			
		Number of leisure-time activities (%)				Number of leisure-time activities (%)			
		0	1	2-4	≥5	0	1	2-4	≥5
Number at risk, n*		6,503 (28.2)	3,848 (16.6)	9,429 (40.8)	3,323 (14.4)	9,071 (33.4)	4,591 (16.9)	10,491 (38.6)	3,030 (11.2)
Age, years**		74.2 (6.4)	74.7 (6.3)	73.6 (5.7)	73.0 (5.3)	75.1 (6.7)	75.2 (6.3)	73.5 (5.7)	72.3 (5.0)
Educational level, years**	≥10	1,797 (27.6)	1,000 (26.0)	3,270 (34.7)	1,201 (36.1)	2,409 (26.6)	1,318 (28.7)	3,961 (37.8)	1,363 (45.0)
Occupational status**	Not employed	3,886 (59.8)	2,387 (62.0)	6,313 (67.0)	2,422 (72.9)	5,586 (61.6)	2,857 (62.2)	7,281 (69.4)	2,343 (77.3)
Equivalized income, 10,000 JPY*	<200	1,395 (21.5)	832 (21.6)	1,388 (14.7)	309 (9.3)	2,012 (22.2)	990 (21.6)	1,662 (15.8)	329 (10.9)
	200-399	3,483 (53.6)	1,980 (51.5)	5,559 (59.0)	2,082 (62.7)	3,883 (42.8)	1,899 (41.4)	5,114 (48.8)	1,596 (52.7)
	≥400	652 (10.0)	425 (11.0)	1,516 (16.1)	728 (21.9)	712 (7.9)	434 (9.5)	1,561 (14.9)	664 (21.9)
Marital status*	Married	5,416 (83.3)	3,187 (82.8)	8,170 (86.7)	2,991 (90.1)	4,892 (53.9)	2,501 (54.5)	6,299 (60.0)	1,904 (62.8)
	Widowed	589 (9.1)	370 (9.6)	776 (8.2)	223 (6.7)	3,265 (36.0)	1,626 (35.4)	3,318 (31.6)	931 (30.7)
	Divorced	201 (3.1)	122 (3.2)	209 (2.2)	61 (1.8)	389 (4.3)	168 (3.7)	397 (3.8)	92 (3.0)
	Never married	131 (2.0)	63 (1.6)	117 (1.2)	23 (0.7)	190 (2.1)	113 (2.5)	270 (2.6)	79 (2.6)
Smoking status*	Current	1,417 (21.8)	834 (21.7)	1,658 (17.6)	417 (12.6)	383 (4.2)	164 (3.6)	248 (2.4)	55 (1.8)
	Former	2,405 (37.0)	1,409 (36.6)	4,161 (44.1)	1,693 (51.0)	334 (3.7)	154 (3.4)	369 (3.5)	97 (3.2)

	Never	1,488 (22.9)	954 (24.8)	2,237 (23.7)	826 (24.9)	6,860 (75.6)	3,529 (76.9)	8,661 (82.6)	2,621 (86.5)
Instrumental activities of daily living (IADL) status**	Independent; 5 points out of 5	3,597 (55.3)	2,254 (58.6)	6,626 (70.3)	2,724 (82.0)	6,274 (69.2)	3,474 (75.7)	9,180 (87.5)	2,892 (95.5)
	Not-independent; less than 5points	2,594 (39.9)	1,412 (36.7)	2,415 (25.6)	497 (15.0)	2,479 (27.3)	956 (20.8)	966 (9.2)	70 (2.3)
Walking hours, min/day*	<30	2,704 (41.6)	1,534 (39.9)	2,666 (28.3)	524 (15.8)	3,726 (41.1)	1,771 (38.6)	3,126 (29.8)	565 (18.7)
	30–59	1,884 (29.0)	1,124 (29.2)	3,368 (35.7)	1,303 (39.2)	2,616 (28.8)	1,437 (31.3)	3,743 (35.7)	1,176 (38.8)
	60–89	802 (12.3)	509 (13.2)	1,668 (17.7)	745 (22.4)	1,017 (11.2)	527 (11.5)	1,647 (15.7)	610 (20.1)
	≥90	920 (14.2)	585 (15.2)	1,626 (17.2)	730 (22.0)	1,104 (12.2)	563 (12.3)	1,602 (15.3)	617 (20.4)
History of diabetes mellitus	Yes	980 (15.1)	582 (15.1)	1,415 (15.0)	491 (14.8)	1,048 (11.6)	466 (10.2)	968 (9.2)	264 (8.7)
History of hypertension	Yes	2,429 (37.4)	1,446 (37.6)	3,594 (38.1)	1,245 (37.5)	3,775 (41.6)	1,931 (42.1)	4,255 (40.6)	1,086 (35.8)
History of stroke	Yes	160 (2.5)	90 (2.3)	162 (1.7)	59 (1.8)	85 (0.9)	31 (0.7)	59 (0.6)	19 (0.6)
History of heart disease	Yes	981 (15.1)	570 (14.8)	1,353 (14.4)	460 (13.8)	994 (11.0)	464 (10.1)	907 (8.7)	244 (8.1)
History of mental disorder**	Yes	81 (1.3)	33 (0.9)	53 (0.6)	23 (0.7)	138 (1.5)	60 (1.3)	90 (0.9)	13 (0.4)
History of hearing loss**	Yes	564 (8.7)	325 (8.5)	701 (7.4)	209 (6.3)	731 (8.1)	387 (8.4)	691 (6.6)	149 (4.9)
Political groups/organizations**	Yes	130 (2.0)	153 (4.0)	398 (4.2)	178 (5.4)	103 (1.1)	214 (4.7)	482 (4.6)	136 (4.5)
Volunteer groups**	Yes	239 (3.7)	188 (4.9)	971 (10.3)	657 (19.8)	204 (2.3)	205 (4.5)	1,003 (9.6)	644 (21.3)
Sports groups/clubs**	Yes	131 (2.0)	465 (12.1)	2,249 (23.9)	1,370 (41.2)	115 (1.3)	513 (11.2)	2,859 (27.3)	1,486 (49.0)

Neighborhood associations**	Yes	497 (7.6)	350 (9.1)	1,307 (13.9)	704 (21.2)	306 (3.4)	256 (5.6)	935 (8.9)	447 (14.8)
Senior clubs**	Yes	322 (5.0)	313 (8.1)	1,206 (12.8)	579 (17.4)	509 (5.6)	531 (11.6)	1,758 (16.8)	672 (22.2)
Frequency of meeting friends*	≥4 times/week	667 (10.3)	486 (12.6)	1,212 (12.9)	492 (14.8)	1,233 (13.5)	665 (14.5)	1,629 (15.5)	584 (19.3)
	2-3 times/week	791 (12.2)	615 (16.0)	1,937 (20.5)	833 (25.1)	1,676 (18.5)	1,082 (23.6)	3,221 (30.7)	1,197 (39.5)
	1 time/week	772 (11.9)	590 (20.5)	1,589 (16.9)	634 (19.1)	1,310 (14.4)	842 (18.3)	2,099 (20.0)	582 (19.2)
	1-2 times/month	1,195 (18.4)	700 (18.8)	2,023 (21.5)	744 (22.4)	1,705 (18.8)	909 (19.8)	1,964 (18.7)	419 (13.8)
	less than 1 times/year	2,734 (42.1)	1,286 (33.5)	2,441 (25.9)	123 (17.4)	2,447 (27.0)	797 (17.4)	1,161 (11.1)	169 (5.6)

All data are presented by means (standard deviations) or numbers (percentages).

**P* for differences in proportions across the number of leisure-time activities: *P*<0.001.

** *P* for trends in means and proportions across the number of leisure-time activities: *P*<0.001.

leisure-time activities

	Men	Women
	Number (%)	Number (%)
Higher-loading leisure-time activities		
Walking/Running	6,119 (55.4)	4,936 (44.6)
Gardening	5,675 (42.6)	7,653 (57.4)
Traveling	5,354 (48.3)	5,742 (51.7)
Cultivation of agricultural crops	3,118 (46.6)	3,572 (53.4)
Karaoke	2,660 (48.5)	2,826 (48.5)
Fishing	2,488 (96.2)	99 (3.8)
Golf	2,381 (90.0)	264 (10.0)
Photography	2,170 (77.5)	629 (22.5)
Grand golf	2,162 (53.5)	1,875 (46.5)
Exercise/Tai Chi	827 (22.6)	2,829 (77.4)
Mountain climbing	654 (65.1)	351 (34.9)
Instrument performance	465 (30.9)	1,040 (69.1)
Gate ball	370 (50.7)	359 (49.3)
Dance	312 (15.4)	1,710 (84.6)
Chorus/ Folk song	217 (15.2)	1,207 (84.8)
Lower-loading leisure-time activities		
Reading	3,865 (50.4)	3,805 (49.6)
Computer	3,148 (75.9)	1,002 (24.1)
Igo/Shogi/Mahjong	2,513 (91.8)	225 (8.2)
Pachinko	1,511 (74.3)	523 (25.7)
Painting/Picture letter	647 (33.1)	1,309 (66.9)
Calligraphy	573 (31.1)	1,270 (68.9)
Haiku/Tanka/Senryu	416 (37.6)	766 (62.4)
Handicrafts	306 (7.5)	3,782 (92.5)
Tea ceremony/Flower arrangement	84 (6.4)	1,223 (93.6)
Other non-specified activities	2,512 (43.5)	3,262 (56.5)

Table 3. Sex-specific hazard ratio (HR) and 95% confidence interval (CI) for functional disability according to the number of leisure-time activities

	Men				Women			
	Number of leisure-time activities				Number of leisure-time activities			
	0	1	2–4	≥5	0	1	2–4	≥5
Number at risk, n	6,503	3,848	9,429	3,323	9,071	4,591	10,491	3,030
Person-years	30,604	18,469	48,279	17,464	42,907	22,624	54,346	16,240
Number of cases, n	1,643	933	1,522	399	2,712	1,189	1,865	368
Age-adjusted HRs (95% CIs)	Ref	0.86 (0.79–0.93)	0.61 (0.57–0.65)	0.47 (0.42–0.52)	Ref	0.78 (0.73–0.84)	0.65 (0.61–0.69)	0.52 (0.47–0.58)
Multivariable HRs ^{1*} (95% CIs)	Ref	0.87 (0.80–0.94)	0.69 (0.64–0.74)	0.60 (0.54–0.67)	Ref	0.82 (0.77–0.88)	0.74 (0.70–0.79)	0.65 (0.58–0.72)
Multivariable HRs ^{2**} (95% CIs)	Ref	0.89 (0.82–0.97)	0.72 (0.67–0.78)	0.66 (0.58–0.74)	Ref	0.84 (0.78–0.90)	0.77 (0.72–0.82)	0.70 (0.62–0.79)
Onset within 1 year excluded								
Number at risk, n	5,159	3,056	8,075	2,971	6,882	3,570	8,870	2,707
Person-years	26,194	15,818	43,532	16,197	35,730	19,124	48,741	15,081
Number of cases, n	1,344	792	1,354	352	2,189	1,021	1,621	323
Age-adjusted HRs (95% CIs)	Ref	0.89 (0.81–0.97)	0.65 (0.60–0.70)	0.49 (0.44–0.55)	Ref	0.82 (0.76–0.88)	0.68 (0.63–0.72)	0.54 (0.48–0.61)
Multivariable HRs ^{1*} (95% CIs)	Ref	0.89 (0.82–0.98)	0.72 (0.67–0.78)	0.61 (0.54–0.69)	Ref	0.86 (0.80–0.92)	0.77 (0.72–0.82)	0.66 (0.58–0.75)
Multivariable HRs ^{2**} (95% CIs)	Ref	0.91 (0.84–0.99)	0.75 (0.69–0.81)	0.66 (0.58–0.75)	Ref	0.87 (0.81–0.94)	0.79 (0.73–0.85)	0.70 (0.62–0.80)
Onset within 2 year excluded								
Number at risk, n	5,486	3,234	8,326	3,026	7,387	3,780	9,186	2,762
Person-years	26,693	16,082	43,913	16,277	36,480	19,436	49,217	15,164
Number of cases, n	1,017	614	1,103	297	1,684	811	1,305	268
Age-adjusted HRs (95% CIs)	Ref	0.90 (0.82–0.99)	0.68 (0.63–0.74)	0.53 (0.46–0.60)	Ref	0.83 (0.76–0.90)	0.69 (0.64–0.74)	0.56 (0.49–0.64)

Multivariable HRs ^{1*} (95% CIs)	Ref	0.91 (0.82–1.00)	0.76 (0.69–0.83)	0.65 (0.57–0.75)	Ref	0.87 (0.80–0.94)	0.76 (0.71–0.82)	0.66 (0.58–0.75)
Multivariable HRs ^{2**} (95% CIs)	Ref	0.93 (0.84–1.02)	0.79 (0.72–0.87)	0.70 (0.61–0.81)	Ref	0.88 (0.81–0.96)	0.79 (0.73–0.85)	0.71 (0.61–0.81)
Onset within 3 year excluded								
Number at risk, n	5,767	3,385	8,597	3,099	7,880	3,985	9,537	2,820
Person-years	27,394	16,460	44,584	16,460	37,699	19,944	50,094	15,312
Number of cases, n	736	463	832	224	1,191	606	954	210
Age-adjusted HRs (95% CIs)	Ref	0.94 (0.84–1.06)	0.70 (0.63–0.77)	0.54 (0.46–0.62)	Ref	0.87 (0.79–0.96)	0.69 (0.63–0.75)	0.59 (0.51–0.69)
Multivariable HRs ^{1*} (95% CIs)	Ref	0.95 (0.85–1.07)	0.78 (0.70–0.86)	0.67 (0.57–0.78)	Ref	0.90 (0.82–0.99)	0.77 (0.70–0.83)	0.69 (0.59–0.80)
Multivariable HRs ^{2**} (95% CIs)	Ref	0.97 (0.87–1.09)	0.81 (0.73–0.90)	0.71 (0.60–0.83)	Ref	0.91 (0.82–1.00)	0.77 (0.71–0.85)	0.72 (0.61–0.85)

* Multivariable HRs¹ were adjusted for age, educational level, occupational status, equivalized income, marital status, smoking status, instrumental activities of daily living (IADL) status, walking hours, histories of diabetes mellitus, hypertension, stroke, heart disease, mental disorder, and hearing loss.

** Multivariable HRs² were adjusted further for sports groups/clubs, senior clubs, neighborhood associations, volunteer groups, political groups/organizations, and frequency of meeting friends.

All of the *P* for trend were <0.001.

Table 4. Hazard ratio (HR) and 95% confidence intervals (CI) for functional disability according to the number of higher- and lower- loading leisure-time activities

	Number of higher-loading leisure-time activities				Number of lower-loading leisure-time activities			
	0	1	2–4	≥5	0	1	2–4	≥5
Number at risk, n	19,759	10,561	17,680	2,196	31,987	11,833	6,381	85
Person-years	93,310	52,983	92,813	11,828	157,643	60,180	32,661	450
Number of cases, n	5,460	2,332	2,597	242	7,489	2,111	1,017	14
Age-adjusted HRs (95% CIs)	Ref	0.76 (0.73–0.80)	0.57 (0.54–0.59)	0.46 (0.41–0.53)	Ref	0.81 (0.77–0.85)	0.74 (0.70–0.80)	0.74 (0.44–1.24)
Multivariable HRs (95% CIs)	Ref	0.83 (0.79–0.87)	0.70 (0.66–0.73)	0.66 (0.58–0.76)	Ref	0.89 (0.85–0.94)	0.88 (0.82–0.94)	1.02 (0.60–1.72)
Onset within 1 year excluded								
Number at risk, n	15,291	8,654	15,360	1,985	25,714	9,998	5,506	72
Person-years	78,648	46,107	84,604	11,058	136,444	53,916	29,661	397
Number of cases, n	4,468	1,997	2,320	211	6,273	1,835	875	13
Age-adjusted HRs (95% CIs)	Ref	0.78 (0.74–0.83)	0.60 (0.57–0.63)	0.47 (0.41–0.54)	Ref	0.83 (0.79–0.87)	0.75 (0.70–0.81)	0.80 (0.47–1.38)
Multivariable HRs (95% CIs)	Ref	0.84 (0.80–0.89)	0.71 (0.68–0.75)	0.65 (0.56–0.75)	Ref	0.91 (0.86–0.96)	0.88 (0.82–0.95)	1.07 (0.62–1.85)
Onset within 2 year excluded								
Number at risk, n	16,345	9,057	15,767	2,018	27,060	10,381	5,674	72
Person-years	80,219	46,717	85,219	11,108	138,476	54,481	29,910	397
Number of cases, n	3,414	1,594	1,913	178	4,927	1,452	707	13
Age-adjusted HRs (95% CIs)	Ref	0.81 (0.76–0.86)	0.62 (0.59–0.66)	0.50 (0.43–0.58)	Ref	0.83 (0.78–0.88)	0.76 (0.71–0.83)	1.01 (0.59–1.74)
Multivariable HRs (95% CIs)	Ref	0.86 (0.81–0.91)	0.74 (0.69–0.79)	0.67 (0.57–0.78)	Ref	0.91 (0.85–0.96)	0.89 (0.82–0.96)	1.31 (0.76–2.26)
Onset within 3 year								

excluded

Number at risk, n	17,296	9,466	16,248	2,060	28,385	10,762	5,848	75
Person-years	82,582	47,732	86,422	11,212	141,760	55,440	30,344	404
Number of cases, n	2,463	1,185	1,432	136	3,602	1,071	533	10
Age-adjusted HRs (95% CIs)	Ref	0.82 (0.77–0.88)	0.63 (0.59–0.67)	0.50 (0.42–0.60)	Ref	0.83 (0.78–0.89)	0.78 (0.72–0.86)	1.08 (0.58–2.00)
Multivariable HRs (95% CIs)	Ref	0.87 (0.81–0.93)	0.73 (0.68–0.78)	0.65 (0.54–0.78)	Ref	0.91 (0.85–0.97)	0.91 (0.83–1.00)	1.38 (0.74–2.57)

Multivariable HRs were adjusted for sex, age, educational level, occupational status, equivalized income, marital status, smoking status, instrumental activities of daily living (IADL), walking hours, histories of diabetes mellitus, hypertension, stroke, heart disease, mental disorder, and hearing loss, sports groups/clubs, senior clubs, neighborhood associations, volunteer groups, political groups/organizations, frequency of meeting friends.

All of the *P* for trend were <0.001.

Supplementary Table 1. Sex-specific multivariable hazard ratio (HR) and 95% confidence interval (CI) for functional disability according to the number of higher- and lower- loading leisure-time activities

		Men	Women
Higher-loading leisure-time activities			
Walking/Running			
	Number at risk, n	6,119	4,936
	Person-years	31,672	25,991
	Number of cases, n	813	710
	Multivariable HRs (95% CIs)	0.87 (0.80–0.95) [†]	0.95 (0.87–1.03)
Gardening			
	Number at risk, n	5,675	7,653
	Person-years	29,261	40,096
	Number of cases, n	928	1,304
	Multivariable HRs (95% CIs)	0.95 (0.88–1.02)	0.92 (0.86–0.99) [†]
Traveling			
	Number at risk, n	5,354	5,742
	Person-years	27,993	30,663
	Number of cases, n	682	698
	Multivariable HRs (95% CIs)	0.82 (0.75–0.89) [†]	0.79 (0.73–0.86) [†]
Cultivation of agricultural crops			
	Number at risk, n	3,118	3,572
	Person-years	16,517	19,067
	Number of cases, n	464	608
	Multivariable HRs (95% CIs)	0.86 (0.78–0.95) [†]	0.81 (0.74–0.89) [†]
Karaoke			
	Number at risk, n	2,660	2,826
	Person-years	13,558	14,606
	Number of cases, n	431	576
	Multivariable HRs (95% CIs)	0.97 (0.87–1.07)	1.18 (1.07–1.29)
Fishing			
	Number at risk, n	2,488	99
	Person-years	13,268	519
	Number of cases, n	298	14
	Multivariable HRs (95% CIs)	0.80 (0.71–0.90) [†]	1.25 (0.74–2.12)
Golf			
	Number at risk, n	2,381	264
	Person-years	12,616	1,382
	Number of cases, n	196	24
	Multivariable HRs (95% CIs)	0.69 (0.59–0.81) [†]	0.91 (0.61–1.37)
Photography			
	Number at risk, n	2,170	629
	Person-years	11,145	3,296

	Number of cases, n	320	86
	Multivariable HRs (95% CIs)	0.98 (0.83–1.07)	1.00 (0.83–1.32)
Grand golf			
	Number at risk, n	2,162	1,875
	Person-years	11,244	10,037
	Number of cases, n	378	283
	Multivariable HRs (95% CIs)	0.94 (0.83–1.07)	0.92 (0.80–1.05)
Exercise/Tai chi			
	Number at risk, n	827	2,829
	Person-years	4,215	15,063
	Number of cases, n	123	396
	Multivariable HRs (95% CIs)	1.00 (0.83–1.20)	0.97 (0.87–1.09)
Mountain climbing			
	Number at risk, n	654	351
	Person-years	3,551	1,945
	Number of cases, n	49	24
	Multivariable HRs (95% CIs)	0.72 (0.54–0.96) [†]	0.83 (0.55–1.24)
Instrument performance			
	Number at risk, n	465	1,040
	Person-years	2,403	5,536
	Number of cases, n	59	153
	Multivariable HRs (95% CIs)	0.77 (0.59–1.00)	0.82 (0.70–0.97) [†]
Gate ball			
	Number at risk, n	370	359
	Person-years	1,734	1,805
	Number of cases, n	103	104
	Multivariable HRs (95% CIs)	1.09 (0.89–1.34)	1.02 (0.83–1.25)
Dance			
	Number at risk, n	312	1,710
	Person-years	1,655	9,180
	Number of cases, n	47	231
	Multivariable HRs (95% CIs)	0.93 (0.70–1.25)	0.80 (0.70–0.91) [†]
Chorus/Folk song			
	Number at risk, n	217	1,207
	Person-years	1,093	6,361
	Number of cases, n	40	199
	Multivariable HRs (95% CIs)	1.11 (0.81–1.52)	0.99 (0.86–1.15)
Lower-loading leisure-time activities			
Reading			
	Number at risk, n	3,865	3,805
	Person-years	19,182	19,293
	Number of cases, n	709	697
	Multivariable HRs (95% CIs)	1.14 (1.04–1.24)	1.07 (0.98–1.17)

	Number at risk, n	5,148	1,882
	Person-years	16,556	5,339
	Number of cases, n	338	105
	Multivariable HRs (95% CIs)	0.84 (0.75–0.95) [†]	0.93 (0.76–1.14)
Igo/Shogi/Mahjong			
	Number at risk, n	2,513	225
	Person-years	12,616	1,138
	Number of cases, n	434	30
	Multivariable HRs (95% CIs)	1.03 (0.93–1.14)	0.74 (0.52–1.07)
Pachinko			
	Number at risk, n	1,511	523
	Person-years	7,869	2,770
	Number of cases, n	204	75
	Multivariable HRs (95% CIs)	0.94 (0.81–1.08)	0.86 (0.68–1.08)
Painting/Picture letter			
	Number at risk, n	647	1,309
	Person-years	3,214	6,772
	Number of cases, n	127	210
	Multivariable HRs (95% CIs)	1.02 (0.85–1.22)	0.92 (0.80–1.06)
Calligraphy			
	Number at risk, n	573	1,270
	Person-years	2,733	6,479
	Number of cases, n	141	247
	Multivariable HRs (95% CIs)	1.12 (0.94–1.33)	1.01 (0.89–1.15)
Haiku/Tanka/Senryu			
	Number at risk, n	461	766
	Person-years	2,258	3,696
	Number of cases, n	94	206
	Multivariable HRs (95% CIs)	0.82 (0.66–1.01)	1.20 (1.04–1.38)
Handicrafts			
	Number at risk, n	306	3,782
	Person-years	1,505	19,642
	Number of cases, n	62	617
	Multivariable HRs (95% CIs)	1.04 (0.81–1.34)	0.86 (0.79–0.94) [†]
Tea ceremony/Flower arrangement			
	Number at risk, n	84	1,223
	Person-years	407	6,388
	Number of cases, n	18	222
	Multivariable HRs (95% CIs)	1.38 (0.86–2.20)	1.02 (0.89–1.17)

Multivariable HRs were adjusted for age, educational level, occupational status, equivalized income, marital status, smoking status, instrumental activities of daily living (IADL), walking hours, histories of diabetes

mellitus, hypertension, stroke, heart disease, mental disorder, hearing loss, sports groups/clubs, senior clubs, neighborhood associations, volunteer groups, political groups/organizations, and frequency of meeting friends.

† *P* value were <0.01.

‡ *P* value were <0.05.

- Leisure-time activities had a health benefit in older people.
- Leisure-time activities were associated with a lower risk of functional disability.
- Even lower-loading activities, tended to be associated with the lower risk.
- The findings suggest the importance to engage in various leisure-time activities.

Yoshihiro Fuji: Conceptualization, Methodology, Data curation, Roles/Writing – original draft, Visualization, Formal analysis, Software

Ryoto Sakaniwa: Conceptualization, Supervision, Validation, Formal analysis, Software, Writing – review & editing

Kokoro Shirai: Conceptualization, Supervision, Validation, Formal analysis, Software, Writing – review & editing

Tami Saito: Conceptualization, Validation, Formal analysis, Writing – review & editing

Shigekazu Ukawa: Conceptualization, Validation, Formal analysis, Writing – review & editing

Hiroyasu Iso: Conceptualization, Supervision, Validation, Formal analysis, Software, Writing – review & editing

Katsunori Kondo: Conceptualization, Investigation, Validation, Formal analysis, Funding acquisition, Project Administration, Resources, Writing – review & editing